

UK/TP 98-14
 hep-ph/9809479
 September 1998

ISOSPIN VIOLATION IN HADRONIC B-DECAYS ^a

S. GARDNER

*Department of Physics and Astronomy, University of Kentucky,
 Lexington, KY 40506-0055 USA
 E-mail: svg@ratina.pa.uky.edu*

I present two disparate examples of isospin violation in hadronic B-decays. In the first, the presence of ρ^0 - ω mixing in the decay $B^\mp \rightarrow \rho^\mp \rho^0(\omega) \rightarrow \rho^\mp \pi^+ \pi^-$ permits the extraction of $\text{sgn}(\sin \alpha)$, where α is the usual angle of the unitarity triangle, with minimal hadronic uncertainty. In the second, the presence of π^0 - η, η' mixing can obscure the extraction of $\sin 2\alpha$ from an isospin analysis in $B \rightarrow \pi\pi$ decays.

1 Introduction

In the standard model, CP violation is characterized by a single phase in the Cabibbo-Kobayashi-Maskawa (CKM) matrix, rendering its elements complex. The CKM matrix of the standard model is unitary, so that determining whether or not this is empirically so is a central test of the standard model's veracity¹. Thus, determining whether the angles of the unitarity triangle, α , β , and γ , empirically sum to π and whether its angles are compatible with the measured lengths of its sides lie at the heart of these tests of the standard model.

CP-violating effects in hadronic B-decays will play a crucial role in the determination of α , β , and γ , and many clever methods have been devised to evade the uncertainties the strong interaction would weigh on their extraction¹. Irrespective of the efficacy of these methods, discrete ambiguities in the angles remain, for experiments in the neutral B sector which would measure an unitarity triangle angle, ϕ , determine $\sin 2\phi$ and not ϕ itself². Nevertheless, removing discrete ambiguities is important, for standard model unitarity requires merely $\alpha + \beta + \gamma = \pi$, mod 2π . Determining the precise equality yields another standard model test, for consistency with the measured value of ϵ and the computed B_K parameter suggest that it ought be π ³.

The isospin-violating effects to be discussed here impact the extraction of α , where $\alpha \equiv \arg[-V_{td}V_{tb}^*/(V_{ud}V_{ub}^*)]$ and V_{ij} is an element of the CKM matrix⁴. The first exploits isospin violation to extract $\text{sgn}(\sin \alpha)$ from the rate asymmetry in $B^\mp \rightarrow \rho^\mp \rho^0(\omega) \rightarrow \rho^\mp \pi^+ \pi^-$, where $\rho^0(\omega)$ denotes the ρ^0 - ω interference region^{5,6}, with minimal hadronic uncertainty, removing the

^aTo appear in the proceedings of the *Workshop on CP Violation*, Adelaide, Australia, July 3-8 1998.

$\text{mod}(\pi)$ ambiguity in α consequent to a $\sin 2\alpha$ measurement. An asymmetry emerges only if both a weak and strong phase difference exists between two interfering amplitudes⁷. The strong phase difference is typically small and uncertain, yet in the decay $B^\mp \rightarrow \rho^\mp \rho^0(\omega) \rightarrow \rho^\mp \pi^+ \pi^-$, the presence of the ω resonance not only enhances the asymmetry to some 20%^{5,6} but also permits the determination of the strong phase from $e^+e^- \rightarrow \pi^+ \pi^-$ data for $\pi^+ \pi^-$ invariant masses in the vicinity of the ω resonance⁶.

The second topic concerns the impact of isospin violation on the extraction of $\sin 2\alpha$ from an isospin analysis in $B \rightarrow \pi\pi$ decays⁸. Isospin is broken not only by electroweak effects but also by the u and d quark mass difference. The latter drives $\pi^0 - \eta, \eta'$ mixing⁹, which, in turn, generates an amplitude in $B \rightarrow \pi\pi$ not included in the isospin analysis. Thus, although the effect of electroweak penguins is estimated to be small^{10,11,12}, when all the effects of isospin violation are included, the error in the extracted value of $\sin 2\alpha$ can be significant¹³.

2 ρ^0 - ω Mixing and CP Violation in $B^\mp \rightarrow \rho^\mp \rho^0(\omega) \rightarrow \rho^\mp \pi^+ \pi^-$

Here we consider the extraction of $\text{sgn}(\sin \alpha)$ from $B^\mp \rightarrow \rho^\mp \rho^0(\omega) \rightarrow \rho^\mp \pi^+ \pi^-$, as proposed by Enomoto and Tanabashi⁵. In this channel, the rate asymmetry, which is CP-violating, is also isospin forbidden. If isospin were a perfect symmetry, then the Bose symmetry of the $J = 0$ $\rho^\pm \rho^0$ final state would force it to have isospin $I = 2$. The strong penguin amplitude is also purely $\Delta I = 1/2$, so that no CP violation is possible in this limit. If isospin violating effects are included, however, two effects occur. The penguin operators then possess both $\Delta I = 1/2$ and $\Delta I = 3/2$ character; the latter are generated by electroweak penguin operators and by the isospin-violating effects which distinguish the $\rho^\pm$ from the ρ^0 . Yet isospin violation also generates ρ^0 - ω mixing, so that a $I = 1$ final state is also possible. In our detailed numerical estimates⁶, we find that the strong phase in the ρ^0 - ω interference region is driven by ρ^0 - ω mixing. The sign and magnitude of ρ^0 - ω mixing is fixed by $e^+e^- \rightarrow \pi^+ \pi^-$ data^{14,15}, so that we are able to interpret direct CP violation in this channel to extract $\text{sgn}(\sin \alpha)$.

Resonances can play a strategic role in direct CP violation, for their mass and width can be used to constrain the strong phase and their interference can significantly enhance the CP-violating asymmetry^{16,17}. To see how these effects are realized here, consider the amplitude A for $B^- \rightarrow \rho^- \pi^+ \pi^-$ decay: $A = \langle \pi^+ \pi^- \rho^- | \mathcal{H}^T | B^- \rangle + \langle \pi^+ \pi^- \rho^- | \mathcal{H}^P | B^- \rangle$, where A is given by the sum of the amplitudes corresponding to the tree and penguin diagrams, respectively. Defining the strong phase δ , the weak phase ϕ , and the magnitude r via $A =$

$\langle \pi^+ \pi^- \rho^- | \mathcal{H}^T | B^- \rangle [1 + r e^{i\delta} e^{i\phi}]$ the CP-violating asymmetry A_{CP} is

$$A_{\text{CP}} \equiv \frac{|A|^2 - |\bar{A}|^2}{|A|^2 + |\bar{A}|^2} = \frac{-2r \sin \delta \sin \phi}{1 + 2r \cos \delta \cos \phi + r^2} , \quad (1)$$

where ϕ is $-\alpha$ ⁴. To express δ in terms of the resonance parameters, let t_V be the tree amplitude and p_V be the penguin amplitude to produce a vector meson V . Thus, the tree and penguin amplitudes become

$$\langle \pi^+ \pi^- \rho^- | \mathcal{H}^T | B^- \rangle = \frac{g_\rho}{s_\rho s_\omega} \tilde{\Pi}_{\rho\omega} t_\omega + \frac{g_\rho}{s_\rho} t_\rho , \quad (2)$$

$$\langle \pi^+ \pi^- \rho^- | \mathcal{H}^P | B^- \rangle = \frac{g_\rho}{s_\rho s_\omega} \tilde{\Pi}_{\rho\omega} p_\omega + \frac{g_\rho}{s_\rho} p_\rho , \quad (3)$$

where we have introduced $\tilde{\Pi}_{\rho\omega}$, the effective ρ^0 - ω mixing matrix element¹⁵; g_ρ , the $\rho^0 \rightarrow \pi^+ \pi^-$ coupling; $1/s_V$, the vector meson propagator, with $s_V = s - m_V^2 + im_V \Gamma_V$; and s , the square of the invariant mass of the $\pi^+ \pi^-$ pair. Using

$$r e^{i\delta} e^{i\phi} = \frac{\langle \pi^+ \pi^- \rho^- | \mathcal{H}^P | B^- \rangle}{\langle \pi^+ \pi^- \rho^- | \mathcal{H}^T | B^- \rangle} \quad (4)$$

and⁵:

$$\frac{p_\omega}{t_\rho} \equiv r' e^{i(\delta_q + \phi)} , \quad \frac{t_\omega}{t_\rho} \equiv \tilde{\alpha} e^{i\delta_{\tilde{\alpha}}} , \quad \frac{p_\rho}{p_\omega} \equiv \beta e^{i\delta_\beta} , \quad (5)$$

one finds, to leading order in isospin violation,

$$r e^{i\delta} = \frac{r' e^{i\delta_q}}{s_\omega} \left\{ \tilde{\Pi}_{\rho\omega} + \beta e^{i\delta_\beta} \left(s_\omega - \tilde{\Pi}_{\rho\omega} \tilde{\alpha} e^{i\delta_{\tilde{\alpha}}} \right) \right\} . \quad (6)$$

A $J = 0$, $I = 1$ $\rho^\pm \rho^0$ final state is forbidden by Bose symmetry if isospin is perfect, so that β is non-zero only if electroweak penguin contributions and isospin violation in the $\rho^\pm$ and ρ^0 hadronic form factors are included. Numerically, $|\tilde{\Pi}_{\rho\omega}|/(m_\omega \Gamma_\omega) \gg \beta$ ⁶. Thus, as $s \rightarrow m_\omega^2$,

$$r e^{i\delta} \rightarrow \frac{r' e^{i\delta_q} \tilde{\Pi}_{\rho\omega}}{im_\omega \Gamma_\omega} , \quad (7)$$

so that $\delta \rightarrow \delta_q - \pi/2$ and $r \rightarrow r' \tilde{\Pi}_{\rho\omega}/m_\omega \Gamma_\omega$ in this limit. The asymmetry is thus determined by the resonance parameters $\tilde{\Pi}_{\rho\omega}$, m_ω , and Γ_ω ; the weak phase ϕ ; and the “short distance” parameters r' and δ_q . The latter are calculable within the context of the operator product expansion if the factorization approximation is applied, though a ratio of hadronic form factors enters as




Figure 1: The CP-violating asymmetry, Eq. [1], in percent, plotted versus the invariant mass q of the $\pi^+\pi^-$ pair in MeV for $[N_c, k^2/m_\rho^2]$. The asymmetries with $\tilde{\Pi}_{\rho\omega} = -3500 \text{ MeV}^2$ and $\tilde{\varepsilon} = -0.005$ are shown for $[2, 0.5]$ (solid line), $[3, 0.5]$ (long-dashed), $[2, 0.3]$ (dashed), and $[3, 0.3]$ (dot-dashed).

well. Here that ratio is modified from unity only by isospin-violating effects which distinguish the $\rho^\pm$ from the ρ^0 ; its deviation from unity is no greater than 1%⁶. The presence of ρ^0 - ω mixing implies that the asymmetry as $s \rightarrow m_\omega^2$ depends on $\cos \delta_q$, *not* $\sin \delta_q$, so that its magnitude is insensitive to the value of the small parameter δ_q . The asymmetry for various N_c and k^2 , where k is the momentum of the virtual boson, as a function of s is shown in Fig. 1, where we have used the effective Hamiltonian of Deshpande and He¹⁰, in QCD to next-to-leading logarithmic (NLL) order, in the factorization approximation¹⁸.

To extract the sign of $\sin \alpha$, note that for $s \approx m_\omega^2$ we have

$$r \sin \delta \approx \frac{\tilde{\Pi}_{\rho\omega} r'}{|s_\omega|^2} [(s - m_\omega^2) \sin \delta_q - m_\omega \Gamma_\omega \cos \delta_q] ; \quad (8)$$

the sign of $\sin \delta$ at $s = m_\omega^2$ is that of $-\tilde{\Pi}_{\rho\omega} \cos \delta_q$. The sign and magnitude of $\tilde{\Pi}_{\rho\omega}$ is determined from a fit to the time-like pion form factor, $F_\pi(s)$, as measured in $e^+e^- \rightarrow \pi^+\pi^-$, where¹⁵

$$F_\pi(s) = F_\rho(s) \left[1 + \frac{1}{3} \frac{\tilde{\Pi}_{\rho\omega}(s)}{s - m_\omega^2 + im_\omega \Gamma_\omega} \right] . \quad (9)$$

Fitting $e^+e^- \rightarrow \pi^+\pi^-$ data we find $\tilde{\Pi}_{\rho\omega} = -3500 \pm 300 \text{ MeV}^2$ ¹⁵. The error is statistical only; the value of $\tilde{\Pi}_{\rho\omega}$ is insensitive to the theoretical ambiguities in

the ρ parametrization, $F_\rho(s)$ ¹⁵. Finally, then, the determination of $\text{sgn}(\sin \alpha)$ relies on that of $\text{sgn}(\cos \delta_q)$. The latter cannot be determined without additional theoretical input; however, it is worth noting that, in our computations in the factorization approximation, $\cos \delta_q < 0$ for all N_c and k^2 ⁶. The situation, finally, is not dissimilar from that of the determination of $\text{sgn}(\sin 2\alpha)$ in neutral B decays to CP eigenstates, for fixing the sign of $\sin 2\alpha$ requires that of the parameter B_B ¹⁹. Nevertheless, the computation of $\cos \delta_q$ can be tested through the measurement of the “skew” of the asymmetry in Eqs. (1) and (8) from a Breit-Wigner shape, noting Fig. 1. Indeed, the shape of the asymmetry yields $\tan \delta_q$ and thus offers a sensitive test of factorization⁶. This is possible as other effects which would skew the asymmetry are smaller still. For example, we have assumed $\tilde{\Pi}_{\rho\omega}$ to be both real and s -independent, yet if we include these effects in our fits to $e^+e^- \rightarrow \pi^+\pi^-$ data, the phase and s -dependence of $\tilde{\Pi}_{\rho\omega}$ in the ρ^0 - ω interference region are statistically consistent with zero¹⁵ and, moreover, do not mar our interpretation of the skew⁶.

In summary, the rate asymmetry in $B^\mp \rightarrow \rho^\mp \rho^0(\omega) \rightarrow \rho^\mp \pi^+ \pi^-$ is large and robust with respect to the known strong interaction uncertainties²⁰. The presence of isospin violation in this decay permits the determination of $\text{sgn}(\sin \alpha)$ once the sign of $\cos \delta_q$ is known, noting $|\cos \delta_q| \sim \mathcal{O}(1)$. The latter can be calculated, yet the shape of the asymmetry yields a direct test of the suitability of our estimate. $\tan \delta_q$ can also be extracted from a comparison with the decay $B^\pm \rightarrow \rho^\pm \omega \rightarrow \rho^\pm \pi^+ \pi^0 \pi^-$; here the asymmetry goes as $\sin \delta_q$ ²¹. The sign of $\sin \alpha$ can be thus extracted with minimal hadronic uncertainty, removing the $\text{mod}(\pi)$ ambiguity in α consequent to a $\sin 2\alpha$ measurement.

3 π^0 - η , η' Mixing in $B \rightarrow \pi\pi$ Decays

To review the isospin analysis in $B \rightarrow \pi\pi$ decays, due to Gronau and London⁸, let us consider the time-dependent asymmetry $A(t)$ ⁴:

$$A(t) = \frac{(1 - |r_{f_{CP}}|^2)}{(1 + |r_{f_{CP}}|^2)} \cos(\Delta m t) - \frac{2(\text{Im } r_{f_{CP}})}{(1 + |r_{f_{CP}}|^2)} \sin(\Delta m t), \quad (10)$$

where $r_{f_{CP}} = (V_{tb}^* V_{td} / V_{tb} V_{td}^*)(\bar{A}_{f_{CP}} / A_{f_{CP}}) \equiv e^{-2i\phi_m} \frac{\bar{A}_{f_{CP}}}{A_{f_{CP}}}$, $A_{f_{CP}} \equiv A(B_d^0 \rightarrow f_{CP})$, and $\Delta m \equiv B_H - B_L$. Denoting the amplitudes $B^+ \rightarrow \pi^+ \pi^0$, $B^0 \rightarrow \pi^0 \pi^0$, and $B^0 \rightarrow \pi^+ \pi^-$ by A^{+0} , A^{00} , and A^{+-} , respectively, and introducing A_I to denote an amplitude of final-state isospin I , we have⁸

$$\frac{1}{2} A^{+-} = A_2 - A_0 \quad ; \quad A^{00} = 2A_2 + A_0 \quad ; \quad \frac{1}{\sqrt{2}} A^{+0} = 3A_2, \quad (11)$$

where analogous relations exist for A^{-0} , $\bar{A}^{00}$, and $\bar{A}^{+-}$ in terms of $\bar{A}_2$ and $\bar{A}_0$. If isospin is perfect, the Bose symmetry of the $J = 0$ $\pi\pi$ state permits

amplitudes of $I = 0, 2$, so that the amplitude $B^\pm \rightarrow \pi^\pm \pi^0$ is purely $I = 2$. Moreover, the strong penguin contributions are of $\Delta I = 1/2$ character, so that they cannot contribute to the $I = 2$ amplitude and no CP violation is possible in the $\pi^\pm \pi^0$ final states. This is identical to the situation in $B^\pm \rightarrow \rho^\pm \rho^0$. The penguin contribution in $B^0 \rightarrow \pi^+ \pi^-$, or in $\bar{B}^0 \rightarrow \pi^+ \pi^-$, can then be isolated and removed by determining the relative magnitude and phase of the $I = 0$ to $I = 2$ amplitudes. We have

$$r_{\pi^+ \pi^-} = e^{-2i\phi_m} \frac{(\bar{A}_2 - \bar{A}_0)}{(A_2 - A_0)} = e^{2i\alpha} \frac{(1 - \bar{z})}{(1 - z)}, \quad (12)$$

where $z(\bar{z}) \equiv A_0/A_2(\bar{A}_0/\bar{A}_2)$ and $\bar{A}_2/A_2 = \exp(-2i\phi_t)$ with $\phi_t \equiv \arg(V_{ud}V_{ub}^*)$ and $\phi_m + \phi_t = \beta + \gamma = \pi - \alpha$ in the standard model⁴. Given $|A^{+-}|$, $|A^{00}|$, $|A^{+0}|$, and their charge conjugates, the measurement of $\text{Im } r_{\pi^+ \pi^-}$ determines $\sin 2\alpha$, modulo discrete ambiguities in $\arg((1 - \bar{z})/(1 - z))$. The latter can be removed via a measurement of $\text{Im } r_{\pi^0 \pi^0}$ as well⁸.

We examine the manner in which isospin-violating effects impact this extraction of $\sin 2\alpha$, for isospin is broken not only by electroweak effects but also by the u and d quark mass difference. Both sources of isospin violation generate $\Delta I = 3/2$ penguin contributions, but the latter also generates $\pi^0 - \eta, \eta'$ mixing⁹, admitting an $I = 1$ amplitude. Although electroweak penguins are estimated to be small^{10,11,12}, other isospin-violating effects, such as $\pi^0 - \eta, \eta'$ mixing, can also be important^{13,22}.

To include the effects of $\pi^0 - \eta, \eta'$ mixing, we write the pion mass eigenstate $|\pi^0\rangle$ in terms of the $SU(3)_f$ perfect states $|\phi_3\rangle = |u\bar{u} - d\bar{d}\rangle/\sqrt{2}$, $|\phi_8\rangle = |u\bar{u} + d\bar{d} - 2s\bar{s}\rangle/\sqrt{6}$, and $|\phi_0\rangle = |u\bar{u} + d\bar{d} + s\bar{s}\rangle/\sqrt{3}$. To leading order in isospin violation⁹

$$|\pi^0\rangle = |\phi_3\rangle + \varepsilon(\cos\theta|\phi_8\rangle - \sin\theta|\phi_0\rangle) + \varepsilon'(\sin\theta|\phi_8\rangle + \cos\theta|\phi_0\rangle), \quad (13)$$

where $|\eta\rangle = \cos\theta|\phi_8\rangle - \sin\theta|\phi_0\rangle + O(\varepsilon)$, and $|\eta'\rangle = \sin\theta|\phi_8\rangle + \cos\theta|\phi_0\rangle + O(\varepsilon')$. Expanding QCD to leading order in $1/N_c$, momenta, and quark masses and diagonalizing the quadratic terms in ϕ_3 , ϕ_8 , and ϕ_0 of the resulting effective Lagrangian determines the mass eigenstates π^0 , η , and η' and yields $\varepsilon = \varepsilon_0 \chi \cos\theta$ and $\varepsilon' = -2\varepsilon_0 \tilde{\chi} \sin\theta$, where $\chi = 1 + (4m_K^2 - 3m_\eta^2 - m_\pi^2)/(m_\eta^2 - m_\pi^2) \approx 1.23$, $\tilde{\chi} = 1/\chi$, $\varepsilon_0 \equiv \sqrt{3}(m_d - m_u)/(4(m_s - \hat{m}))$, and $\hat{m} \equiv (m_u + m_d)/2$ ⁹. Thus the magnitude of isospin breaking is controlled by the $SU(3)$ -breaking parameter $m_s - \hat{m}$. The $\eta - \eta'$ mixing angle θ is found to be $\sin 2\theta = -(4\sqrt{2}/3)(m_K^2 - m_\pi^2)/(m_\eta^2 - m_\pi^2) \approx -22^\circ$ ⁹. The resulting $\varepsilon = 1.14\varepsilon_0$ is comparable to the one-loop-order chiral perturbation theory result of $\varepsilon = 1.23\varepsilon_0$ in $\eta \rightarrow \pi^+ \pi^- \pi^0$ ^{23,9}. Using $m_q(\mu = 2.5 \text{ GeV})$ of Ali *et al.*²⁴, we find $\varepsilon = 1.4 \cdot 10^{-2}$ and $\varepsilon' = 7.7 \cdot 10^{-3}$.




Figure 2: Reduced amplitudes in $B \rightarrow \pi\pi$ in the factorization approximation with $[N_c, k^2/m_b^2]$ for a) $[2, 0.5]$, b) $[3, 0.5]$ (solid line) and $[3, 0.3]$ (dashed line), and c) $[\infty, 0.5]$.

In the presence of isospin-violating effects, the $B \rightarrow \pi\pi$ amplitudes become

$$A^{-0} = \langle \pi^- \phi_3 | \mathcal{H}^{\text{eff}} | B^- \rangle + \varepsilon_8 \langle \pi^- \phi_8 | \mathcal{H}^{\text{eff}} | B^- \rangle + \varepsilon_0 \langle \pi^- \phi_0 | \mathcal{H}^{\text{eff}} | B^- \rangle \quad (14)$$

$$\bar{A}^{00} = \langle \phi_3 \phi_3 | \mathcal{H}^{\text{eff}} | \bar{B}^0 \rangle + 2\varepsilon_8 \langle \phi_3 \phi_8 | \mathcal{H}^{\text{eff}} | \bar{B}^0 \rangle + 2\varepsilon_0 \langle \phi_3 \phi_0 | \mathcal{H}^{\text{eff}} | \bar{B}^0 \rangle, \quad (15)$$

where $\varepsilon_8 \equiv \varepsilon \cos \theta + \varepsilon' \sin \theta$ and $\varepsilon_0 \equiv \varepsilon' \cos \theta - \varepsilon \sin \theta$. The $B \rightarrow \pi\pi$ amplitudes satisfy

$$\begin{aligned} \bar{A}^{+-} + 2\bar{A}^{00} - \sqrt{2} A^{-0} &= 4\varepsilon_8 \langle \phi_3 \phi_8 | \mathcal{H}^{\text{eff}} | \bar{B}^0 \rangle + 4\varepsilon_0 \langle \phi_3 \phi_0 | \mathcal{H}^{\text{eff}} | \bar{B}^0 \rangle \\ &\quad - \sqrt{2}\varepsilon_8 \langle \pi^- \phi_8 | \mathcal{H}^{\text{eff}} | B^- \rangle - \sqrt{2}\varepsilon_0 \langle \pi^- \phi_0 | \mathcal{H}^{\text{eff}} | B^- \rangle, \end{aligned} \quad (16)$$

and thus the triangle relation implied by Eq. 11 becomes a quadrilateral. We ignore the small mass differences $m_{\pi^\pm} - m_{\pi^0}$ and $m_{B^\pm} - m_{B^0}$.

We proceed by computing the individual amplitudes using the $\Delta B = 1$ effective Hamiltonian resulting from the operator product expansion in QCD in NLL order²⁴, using the factorization approximation for the hadronic matrix elements. In this context, we can then apply the isospin analysis delineated above to infer $\sin 2\alpha$ and thus estimate its theoretical systematic error, incurred through the neglect of isospin violating effects. Numerical results for

Table 1: Strong phases and inferred values of $\sin 2\alpha$ ⁸ from amplitudes in the factorization approximation with N_c and $k^2/m_b^2 = 0.5$. The strong phase $2\delta_{\text{true}}$ is the opening angle between the $\bar{A}_R^{+-}$ and A_R^{+-} amplitudes in Fig. 1, whereas $2\delta_{\text{GL}}$ is the strong phase associated with the closest matching $\sin 2\alpha$ values, denoted $(\sin 2\alpha)_{\text{GL}}$, from $\text{Im}r_{\pi^+\pi^-}/\text{Im}r_{\pi^0\pi^0}$, respectively. The bounds $|2\delta_{\text{GQI}}|$ and $|2\delta_{\text{GQII}}|$ on $2\delta_{\text{true}}$ from Eqs. 2.12 and 2.15 of Ref. ²⁵ are also shown. All angles are in degrees. We input a) $\sin 2\alpha = 0.0432^{24,26,27}$, b) $\sin 2\alpha = -0.233$ ($\rho = 0.2, \eta = 0.35$)²⁸, and c) $\sin 2\alpha = 0.959$ ($\rho = -0.12$).

case	N_c	$2\delta_{\text{true}}$	$ 2\delta_{\text{GQI}} $	$ 2\delta_{\text{GQII}} $	$ 2\delta_{\text{GL}} $	$(\sin 2\alpha)_{\text{GL}}$
a	2	24.4	26.1	15.8	16.6	-0.0900/-0.0221
a	3	24.2	16.9	16.1	16.2	-0.0926/0.107
a	∞	23.8	59.4	25.1	23.6	0.0451/0.394
b	2	19.6	23.4	12.1	12.9	-0.343/-0.251
b	3	19.4	13.5	12.9	13.0	-0.719/-0.855(*)
b	∞	19.2	59.9	23.6	0.76	-0.550/-0.814(*, †)
c	2	28.3	36.5	20.4	21.0	0.917/0.915
c	3	28.0	24.0	19.1	19.0	0.905/0.952
c	∞	28.3	36.5	20.4	21.0	0.917/0.915

* The matching procedure fails to choose a $\sin 2\alpha$ which is as close to the input value as possible.

† The discrete ambiguity in the strong phase is resolved wrongly.

the reduced amplitudes A_R and $\bar{A}_R$, where $\bar{A}_R^{00} \equiv 2\bar{A}^{00}/((G_F/\sqrt{2})iV_{ub}V_{ud}^*)$, $\bar{A}_R^{+-} \equiv \bar{A}^{+-}/((G_F/\sqrt{2})iV_{ub}V_{ud}^*)$, and $A_R^{-0} \equiv \sqrt{2}A^{-0}/((G_F/\sqrt{2})iV_{ub}V_{ud}^*)$, with $N_c = 2, 3, \infty$ and $k^2/m_b^2 = 0.3, 0.5$ are shown in Fig. 2. A_R^{+0} and A_R^{-0} are broken into tree and penguin contributions, so that $A_R^{+0} \equiv T_{\pi^+\phi_3} + P_{\pi^+\pi^0}$ and $A_R^{-0} \equiv T_{\pi^-\phi_3} + P_{\pi^-\pi^0}$, where $P_{\pi^\pm\pi^0}$ is defined to include the isospin-violating tree contribution in $A_R^{\pm 0}$ as well. The shortest side in each polygon is the vector defined by the RHS of Eq. 16. The values of $\sin 2\alpha$ extracted from the computed amplitudes with N_c and $k^2/m_b^2 = 0.5$ are shown in Table 1 — the results for $k^2/m_b^2 = 0.3$ are similar and have been omitted. In the presence of π^0 - η , η' mixing, the $\bar{A}_R^{+-}$, $\bar{A}_R^{-0}$, and $\bar{A}_R^{00}$ amplitudes obey a quadrilateral relation as per Eq. 16. Consequently, the values of $\sin 2\alpha$ extracted from $\text{Im}r_{\pi^+\pi^-}$ and $\text{Im}r_{\pi^0\pi^0}$ can not only differ markedly from the value of $\sin 2\alpha$ input but also need not match. The incurred error in $\sin 2\alpha$ increases as the value to be extracted decreases; the structure of Eq. 12 suggests this, for as $\sin 2\alpha$ decreases the quantity $\text{Im}((1-\bar{z})/(1-z))$ becomes more important to determining the extracted value. It is useful to contrast the impact of the various isospin-violating effects. The presence of $\Delta I = 3/2$ penguin contributions, be they from $m_u \neq m_d$ or electroweak effects, shift the extracted value of $\sin 2\alpha$ from its input value, yet the “matching” of the $\sin 2\alpha$ values in $\text{Im}r_{\pi^+\pi^-}$ and

$\text{Im}r_{\pi^0\pi^0}$ is unaffected. The mismatch troubles seen in Table 1 are driven by π^0 - η , η' mixing, though the latter shifts the values of $\sin 2\alpha$ in $\text{Im}r_{\pi^+\pi^-}$ as well. Picking the closest matching values of $\sin 2\alpha$ in the two final states also picks the solutions closest to the input value; the exceptions are noted in Table 1. If $|A^{00}|$ and $|\overline{A}^{00}|$ are small⁸ the complete isospin analysis may not be possible, so that we also examine the utility of the bounds recently proposed by Grossman and Quinn²⁵ on the strong phase $2\delta_{\text{true}} \equiv \arg((1 - \bar{z})/(1 - z))$ of Eq. 12. The bounds $2\delta_{\text{GQI}}$ and $2\delta_{\text{GQII}}$ given by their²⁵ Eqs. 2.12 and 2.15, respectively, follow from Eq. 11, and thus can be broken by isospin-violating effects. As shown in Table 1, the bounds typically are broken, and their efficacy does not improve as the value of $\sin 2\alpha$ to be reconstructed grows large.

To conclude, we have considered the role of isospin violation in $B \rightarrow \pi\pi$ decays and have found the effects to be significant. Most particularly, the utility of the isospin analysis in determining $\sin 2\alpha$ strongly depends on the value to be reconstructed. The error in $\sin 2\alpha$ from a $\text{Im}r_{\pi^+\pi^-}$ measurement can be 50% or more for the small values of $\sin 2\alpha$ currently favored by phenomenology^{24,26,27,28}; however, if $\sin 2\alpha$ were near unity, the error would decrease to less than 10%. The effects found arise in part because the penguin contribution in $B^0 \rightarrow \pi^+\pi^-$, e.g., is itself small; we estimate $|P|/|T| < 9\%|V_{tb}V_{td}^*|/|V_{ub}V_{ud}^*|$. Relative to this scale, the impact of π^0 - η , η' mixing is significant. Yet, were the penguin contributions in $B \rightarrow \pi\pi$ larger, the isospin-violating effects considered would still be germane, for not only would the $\Delta I = 3/2$ penguin contributions likely be larger but the $B \rightarrow \pi\eta$ and $B \rightarrow \pi\eta'$ contributions could also be larger as well²⁹. To conclude, we have shown that the presence of π^0 - η , η' mixing breaks the triangle relationship, Eq. 11, usually assumed⁸ and can mask the true value of $\sin 2\alpha$.

Acknowledgements

I am grateful to H. B. O'Connell and A. W. Thomas for their collaboration on issues related to ρ^0 - ω mixing and to the organizers for their invitation to speak at this workshop. This work was supported by the U.S. Department of Energy under DE-FG02-96ER40989 and by the Special Research Centre for the Subatomic Structure of Matter at the University of Adelaide.

References

- [1] R. Fleischer, *Int. J. Mod. Phys. A* **12**, 2459 (1997).
- [2] Y. Grossman and H.R. Quinn, *Phys. Rev. D* **48**, 2139 (1993).
- [3] Y. Nir and H.R. Quinn, *Phys. Rev. D* **42**, 1473 (1990).
- [4] R.M. Barnett *et al.*, *Phys. Rev. D* **54**, 1 (1996).

- [5] R. Enomoto and M. Tanabashi, *Phys. Lett. B* **386**, 413 (1996); note also H.J. Lipkin, *Phys. Lett. B* **357**, 404 (1995).
- [6] S. Gardner, H.B. O'Connell, and A.W. Thomas, *Phys. Rev. Lett.* **80**, 1834 (1998).
- [7] M. Bander, D. Silverman, and A. Soni, *Phys. Rev. Lett.* **43**, 242 (1979).
- [8] M. Gronau and D. London, *Phys. Rev. Lett.* **65**, 3381 (1990). Note $A_{\text{GL}}^{+-} = A^{+-}/\sqrt{2}$ and $A_{\text{GL}}^{\pm 0} = A^{\pm 0}/\sqrt{2}$.
- [9] H. Leutwyler, *Phys. Lett. B* **374**, 181 (1996).
- [10] N.G. Deshpande and X.-G. He, *Phys. Rev. Lett.* **74**, 26 (1995).
- [11] M. Gronau, O.F. Hernández, D. London, and J.L. Rosner, *Phys. Rev. D* **52**, 6374 (1995).
- [12] R. Fleischer, *Phys. Lett. B* **365**, 399 (1996).
- [13] S. Gardner, hep-ph/9806423.
- [14] G.A. Miller, B.M.K. Nefkens, and I. Šlaus, *Phys. Rept.* **194**, 1 (1990); note Fig. 1.6.
- [15] S. Gardner and H.B. O'Connell, *Phys. Rev. D* **57**, 2716 (1998).
- [16] D. Atwood and A. Soni, *Phys. Rev. Lett.* **74**, 220 (1995); *Z. Phys. C* **64**, 241 (1994).
- [17] D. Atwood, G. Eilam, M. Gronau, A. Soni, *Phys. Lett. B* **341**, 372 (1995); G. Eilam, M. Gronau, and R.R. Mendel, *Phys. Rev. Lett.* **74**, 4984 (1995).
- [18] Here we use the CKM parameters of Ali *et al.* ²⁴, so that $\rho = 0.12$, whereas we used $\rho = -0.12$ previously ⁶. The asymmetries differ by $\mathcal{O}(1)\%$.
- [19] Y. Grossman, B. Kayser, and Y. Nir, *Phys. Lett. B* **415**, 90 (1997).
- [20] ρ^0 - ω mixing also generates a large asymmetry in $\Lambda_b \rightarrow n\rho^0(\omega) \rightarrow n\pi^+\pi^-$ and $\Lambda_b \rightarrow \Lambda\rho^0(\omega) \rightarrow \Lambda\pi^+\pi^-$ decays, see X.-H. Guo and A.W. Thomas, hep-ph/9805332.
- [21] S. Gardner, H.B. O'Connell, and A.W. Thomas, in preparation.
- [22] π^0 - η , η' mixing also generates significant isospin-violating corrections to ϵ'/ϵ , see A.J. Buras and J.-M. Gerard, *Phys. Lett. B* **192**, 156 (1987); J.F. Donoghue, E. Golowich, B.R. Holstein, and J. Trampetic, *Phys. Lett. B* **179**, 361 (1986).
- [23] J. Gasser and H. Leutwyler, *Nucl. Phys. B* **250**, 539 (1985).
- [24] A. Ali, G. Kramer, and C.-D. Lü, hep-ph/9804363.
- [25] Y. Grossman and H.R. Quinn, *Phys. Rev. D* **58**, 017504 (1998).
- [26] A. Ali, hep-ph/9801270; A. Ali and D. London, *Nucl. Phys. B Proc. Suppl.* **54A**, 297 (1997); F. Parodi, P. Roudeau, and A. Stocchi, hep-ph/9802289.
- [27] S. Mele, these proceedings.

- [28] D. London, private communication, from an update of Ref.²⁶.
- [29] M. Ciuchini et al., *Nucl. Phys. B* **501**, 271 (1997); **512**, 3 (1998).